

**EDUCATION COMPANY®** 

Edvo-Kit #S-75

## Do Onions, Strawberries and Bananas Have DNA?

#### **Experiment Objective:**

In this experiment, students will learn about the physical nature of DNA by isolating DNA from onions and using the common procedure of DNA spooling.

See page 3 for storage instructions.

Edvo-Kit #

**S-75** 

## **Table of Contents**

	Page
Experiment Components	3
Experiment Requirements	3
Background Information	4
Experiment Procedures	
Experiment Overview	6
Activity One - Spooling DNA from Solution	7
Activity Two - Staining the Spooled DNA	8
Activity Three - Extraction of DNA from Onion Tissue	9
Activity Four - Extraction of DNA from Strawberries or Banana	10
Study Questions	11
Instructor's Guidelines	12
Pre-Lab Preparations	12
Study Questions and Answers	13

Safety Data Sheets can be found on our website: www.edvotek.com/safety-data-sheets

## **EDVO-TECH Service**

## **1.800.EDVOTEK**

Mon. - Fri. 8am-5:30pm EST

#### Please Have the Following Info:

- Product Number & Description
  - Lot Number on Box
  - · Order/Purchase Order #
  - Approx. Purchase Date

Fax: 202.370.1501 • info@edvotek.com • www.edvotek.com



## www.edvotek.com

- Online Catalog
- Order Products
- Experiment Protocols
  - Tech Support
  - Resources!





1.800.EDVOTEK • Fax 202.370.1501 • info@edvotek.com • www.edvotek.com •

## **Experiment Components**

Contents	Check (√)	
<ul><li>DNA Extraction Buffer</li><li>DNA sample in a capped test tube</li></ul>		Experiment #S-75 is designed for 10 groups.
<ul> <li>Transfer pipets</li> <li>Colored Beads (4 colors)</li> </ul>	<u> </u>	Storage: Store entire experiment
<ul><li> Graduated test tubes</li><li> Spooling rods</li><li> Salt packet</li></ul>		in the refrigerator.

## Requirements

- Fresh onion pieces (scallions, also called green onions, are highly recommended)
- Ripened fresh strawberry or banana pieces
- · 20 ml beaker
- Test tubes (13 x 100 mm)
- 70% clear isopropyl alcohol (rubbing alcohol)
- Distilled water
- Ice

All experiment components are intended for educational research only. They are not to be used for diagnostic or drug purposes, nor administered to or consumed by humans or animals.

EDVOTEK and The Biotechnology Education Company are registered trademarks of EDVOTEK, Inc.



## **Background Information**

#### DNA - DEOXYRIBONUCLEASE

Since the 1800s, it has been known that all living organisms are composed of cells. Organisms such as bacteria are single cells, while very complex organisms, such as humans, are composed of billions of many different cells.


In 1868, a Swiss biologist named Friedrich Meischer carried out research which indicated that the nucleus of cells contains a material which he called nucleic acid. But, it wasn't until much later in the 1940's that the nucleic acid, deoxyribonucleic acid (DNA), was recognized as the carrier of genetic information.

DNA plays an important role in two processes. During the process of replication, DNA provides information to copy itself, so genetic information can be passed on from generation to generation of cells. In its second important role,

DNA provides instructions for making proteins, which are vital to the maintenance and function of cells. DNA provides information for the order of amino acids required for making various proteins.

The structure of the DNA molecule was determined by James Watson and Francis Crick in 1953. They determined that DNA was a double helix consisting of two strands. The Watson and Crick model is often described as a spiral ladder. The two strands of DNA are the backbone of the ladder, made of sugar phosphodiester groups. The sugar backbone acts as a support for the rungs of the ladder. These rungs are composed of the chemical bases Adenine, Guanine, Cytosine, and Thymine. The first letters of these bases, A,G,C, and T, are used by scientists to designate the order of the bases within the DNA strands.

The bases are always arranged in pairs. When A occurs on one strand, T occurs on the opposite strand. Similarly, G and C are opposite DNA strands. The bases are held together by weak bonds which are shown as dashed lines in Figure 1.






#### **Background Information**

DNA can be extracted from the nucleus of cells by adding an aqueous buffered extraction solution to cells. The cells are chemically lysed (broken open) and DNA from chromosomes is released. This procedure is known as cell lysis. DNA is soluble in water and cannot be seen, but it is insoluble in alcohol. Purification procedures for nucleic acids usually include precipitation with alcohol in the presence of salt. Since rubbing alcohol (isopropyl alcohol) has a lower density than water, it will form a second layer above the DNA solution. A spooling rod is used to spool the two liquids at the interface of the two phases to separate the DNA from the solution. The DNA will appear as a viscous, clotted mass as it is collected on the spooling rod (Figure 2). The amount of DNA spooled is a consequence of the size of the DNA fragments which are much larger than the small bio-molecules such as amino acids and small carbohydrate sugars.

In this experiment, DNA will be isolated in one of two ways: 1) DNA will be spooled from a buffer solution containing salts; 2) DNA will be extracted from tissues such as onion, banana, or strawberry and spooled from solution. DNA will be observed by the naked eye on the spooling rod.



## **Experiment Overview**

#### **EXPERIMENT OBJECTIVE:**

In this experiment, students will learn about the physical nature of DNA by isolating DNA from onions and using the common procedure of DNA spooling.

#### LABORATORY SAFETY

- 1. Gloves and goggles should be worn routinely as good laboratory practice.
- 2. Exercise extreme caution when working with equipment that is used in conjunction with the heating and/or melting of reagents.
- 3. DO NOT MOUTH PIPET REAGENTS USE PIPET PUMPS.
- 4. Exercise caution when using any electrical equipment in the laboratory.
- 5. Always wash hands thoroughly with soap and water after handling reagents or biological materials in the laboratory.



#### **WORKING HYPOTHESIS**

DNA can be extracted from the cells, and isolated from solution by spooling.

#### **LABORATORY NOTEBOOKS:**

Scientists document everything that happens during an experiment, including experimental conditions, thoughts and observations while conducting the experiment, and, of course, any data collected. Today, you'll be documenting your experiment in a laboratory notebook or on a separate worksheet.

#### Before starting the Experiment:

- Carefully read the introduction and the protocol. Use this information to form a hypothesis for this experiment.
- Predict the results of your experiment.

#### During the Experiment:

Record your observations.

#### After the Experiment:

- Interpret the results does your data support or contradict your hypothesis?
- If you repeated this experiment, what would you change? Revise your hypothesis to reflect this change.



## **Activity One - Construction DNA Models**

#### THE EXPERIMENT PROCEDURE

- 1. Each group of students will get a set of beads of various colors.
- 2. The teacher will assign a base designation for four of the colors. For example:

```
Red = Adenine (A) Blue = Thymine (T)
Green = Guanine (G) Yellow = Cytosine (C)
```

3. Sort all the beads by color. Based upon the beads you received, correctly put the colored beads together in two separate strands to form one or more of the following base pair models:

```
A-T base pair G-C base pair
```

How many A-T or G-C base pairs were you able to put together?

#### PUTTING ALL THE STUDENT BASE PAIR MODELS TOGETHER:

- 4. With the help of your teacher, and the rest of the class, build a model of the DNA ladder. Attach the ends of several bead sections together to form two long strands of double-stranded DNA.
- 7. In your own words, describe the structure of DNA.



## **Activity Two - Teacher Demonstration of DNA Spooling**

In this activity, DNA will be spooled from an aqueous buffered solution by adding salt and over laying with isopropyl alcohol. The isopropyl alcohol (clear rubbing alcohol) should be ice cold (stored in a freezer at least 2 hours before use). Just before the demonstration, place the DNA solution and the alcohol on ice.

#### THE CONTROL

- 1. Add two pinches of salt to a small beaker or clear graduated test tube.
- 2. Add distilled water up to the 2 ml level of the tube and swirl to dissolve the salt.
- 3. Use a transfer pipet to add 1 ml of the control solution (distilled water). Swirl to mix.
- 4. Using a transfer pipet, carefully overlay the control solution with two volumes of very cold 70% clear isopropyl alcohol (about 6 ml). Let the cold alcohol gently flow down the side of the beaker or test tube. Do not mix the two solutions.
- 5. Place the end of the spooling rod just below the line separating the two solutions (the interface). Twirl the rod with two fingers and spool.
- 6. Record your observations for the control.
- 7. Rinse the rod with distilled water and thoroughly dry with a paper towel.

#### THE EXPERIMENT

- 8. Add two pinches of salt to a small beaker or clear graduated test tube.
- 9. Use a transfer pipet to add 2 ml of distilled water and swirl to dissolve the salt.
- 10. Use a transfer pipet to transfer all of the DNA solution to the salt water. Stir or swirl to mix. The total volume should be approximately 3 ml.
- 11. Layer the cold alcohol on top of the DNA solution in the same manner as step #4 of the experiment. Do not mix the two solutions.
- 12. Place the end of the spooling rod just below the line separating the two solutions (the interface). Twirl the rod with two fingers and spool the DNA from the solution.
- 13. Observe the DNA and record your observations.





1.800.EDVOTEK • Fax 202.370.1501 • info@edvotek.con • www.edvotek.con

## **Activity Three - Extraction of DNA from Onion Tissue**


In this activity, DNA is extracted from onion tissue using an extraction buffer and spooled from solution by overlaying the liquid with ice cold isopropyl alcohol.

#### THE EXPERIMENT

- 1. Carefully slice a 5 x 5 x 5 mm cube (about the size of a pencil eraser) from the main body of the onion (not the root tip), and place it in one of the test tubes provided.
- 2. Using a transfer pipet, add DNA Extraction Buffer until it reaches the 3 ml level mark on the tube. Mince the onion cube with the tip of a spooling rod, pencil or pen (mash the onion very well). This will release the cellular contents and the DNA in the onion cells.
- 3. With a transfer pipet, carefully remove 2 ml of the liquid layer from the test tube and transfer it into a second, clean test tube. Try to minimize carry-over of the onion tissue.
- 4. Carefully overlay the liquid with 4 ml of **very cold** 70% clear Isopropyl alcohol.
- 5. Place a spooling rod into the test tube and twirl it with two of your fingers. Gently rotate the rod at the interface of the two phases. The DNA will begin to spool (wrap) around the rod.
  - Gently lift the rod out of the solution from time to time and observe the DNA substance attached to it.
- 6. After spooling for several minutes, remove the rod from the test tube to observe the DNA. The DNA will appear as a viscous material adhering to the splint (it may be grayish in color if a pencil was used to mince the onion tissue).
- 7. Record your observations of the onion DNA.

#### THE CONTROL

8. Based upon what you have previously learned, outline and perform the control for this activity.






## **Activity Four - Extraction of DNA from Strawberries or Banana**

In this activity, DNA is extracted from a banana or strawberries in the same manner as the extraction of DNA from onion.

- 1. Carefully slice a  $5 \times 5 \times 5$  mm cube from the fleshy body of the banana or strawberry, and place it in a clean test tube.
- 2. Using a transfer pipet, add DNA Extraction Buffer until it reaches the 3 ml mark on the tube. Mince the cube with the tip of a spooling rod, pencil or pen to release the cellular contents.



- 3. With a transfer pipet, carefully remove 2 ml of the liquid layer from the test tube and transfer it into a second, clean test tube. Try to minimize carry-over of the banana or strawberry tissue.
- 4. Carefully overlay the liquid with 4 ml of **very cold** 70% clear Isopropyl alcohol.
- 5. Place a spooling rod into the test tube and twirl it with two of your fingers. Gently rotate the rod at the interface of the two phases. The DNA will begin to spool (wrap) around the rod.
  - Gently lift the rod out of the solution from time to time and observe the DNA substance attached to it.
- 6. After spooling for several minutes, remove the rod from the test tube to observe the DNA. The DNA will appear as a viscous material adhering to the splint (it may be grayish in color if a pencil was used to mince the onion tissue).
- 7. Record your observations of the banana or strawberry DNA.



## **Study Questions**

- 1. Describe the appearance of the isolated DNA.
- 2. What is a nucleus?
- 3. What are nucleotides?
- 4. What are chromosomes?
- 5. What does cell lysis mean?
- 6. Why did the alcohol layer on top of the DNA solution?
- 7. Why was it important not to mix the alcohol and DNA solutions?
- 8. What properties of DNA allow spooling?
- 9. What difficulties did you have in spooling? Why?
- 10. How many chromosomes do humans have?

## Instructor's Guide

#### **Pre-Lab Preparations:**

#### ON THE DAY OF THE LAB:

1. Thoroughly chill the 70% isopropyl alcohol (rubbing alcohol) before the students perform the experiment.



- Store it in the freezer for several hours or overnight to ensure that it is very cold.
- Using a calibrated transfer pipet, transfer 6 ml of 70% isopropyl alcohol into 20 test tubes.
- · Place the alcohol-filled tubes back in the freezer or on ice until students are ready to use the alcohol.

### **EDVO-TECH Service**

**1.800.EDVOTEK** 

Mon. - Fri. 8am-5:30pm EST

# 2

#### Please Have the Following Info:

- Product Number & Description
  - · Lot Number on Box
  - Order/Purchase Order #
  - Approx. Purchase Date

Fax: 202.370.1501 • info@edvotek.com • www.edvotek.com

## www.edvotek.com

- Online Catalog
- Order Products
- Experiment Protocols
  - Tech Support
  - Resources!





1.800.EDVOTEK • Fax 202.370.1501 • info@edvotek.com • www.edvotek.com !

## Please refer to the kit insert for the Answers to Study Questions